

## Case 3424

### PSEUDOCOCCINI Cockerell, 1905 (Insecta, Hemiptera): proposed precedence over SPHAEROCOCCINI Cockerell, 1899

Nate B. Hardy and Penny J. Gullan

*Department of Entomology, University of California, One Shields Avenue, Davis, CA 95616–8584, U.S.A. (emails: nbhardy@ucdavis.edu; pjgullan@ucdavis.edu)*

**Abstract.** The purpose of this application, under Article 23.9.3 of the Code, is to conserve the usage of the family-group name PSEUDOCOCCINI Cockerell, 1905 (type genus *Pseudococcus* Westwood, 1840) for the mealybugs, a speciose group including many pests of agriculture, by giving it precedence over the senior family-group name SPHAEROCOCCINI Cockerell, 1899 (type genus *Sphaerococcus* Maskell, 1892).

**Keywords.** Nomenclature; taxonomy; Hemiptera; COCCOIDEA; PSEUDOCOCCINI; SPHAEROCOCCINI; *Pseudococcus*; *Sphaerococcus*; mealybugs.

---

1. Maskell (1892, p. 39) established the generic name *Sphaerococcus*, for the Australian legless mealybug species *S. casuarinae* Maskell, 1892 (p. 39), type species by monotypy.

2. Cockerell (1899, p. 389) placed *Sphaerococcus* and a group of additional legless mealybug genera in a new tribe SPHAEROCOCCINI of the family COCCIDAE Fallén, 1814 (p. 23).

3. Cockerell (1905, p. 193) erected the tribe PSEUDOCOCCINI within the COCCIDAE for *Pseudococcus* Westwood, 1840 (p. 448) and the other mealybug genera known at that time to occur in the Rocky Mountains of North America.

4. Lobdell (1930, p. 209) was the first to use the family name PSEUDOCOCCIDAE for the mealybugs (including SPHAEROCOCCINI) a practice subsequently followed by all entomologists. The name SPHAEROCOCCINI has never been used as a family name and has never been used outside of discussions of mealybug subfamily-level classification.

5. The history of the generic name *Pseudococcus* Westwood (1840, p. 448) is extremely complicated. Miller (1974) (BZN 31: 146–153) submitted a case to the Commission, and in Opinion 1247 (July, 1983, BZN 40: 77–80), the Commission placed *Pseudococcus* Westwood, 1840 (type species *Dactylopius longispinus* Targioni-Tozzetti, 1876) on the Official List of Generic Names in Zoology (Name Number 2188), and PSEUDOCOCCIDAE Cockerell, 1905 (type genus *Pseudococcus* Westwood, 1840) on the Official List of Family-Group Names (Name Number 555).

6. The adoption of SPHAEROCOCCINI in place of its junior synonym PSEUDOCOCCINI would change the name of one of the most commonly encountered pests of agriculture, and leave an enormous body of literature divorced from classification (as of 17 May 2007, 1080 publications in Zoological Record use the name PSEUDOCOCCIDAE). Article 35.5 does not apply in this case, as *P. longispinus* and *S. casuarinae* are in the same subfamily and tribe. We therefore propose that

family-group names based on *Pseudococcus* be given precedence over those based on *Sphaerococcus*. The family-group names based on *Sphaerococcus* would remain available for any coccidologist who may in the future consider the two type genera to belong to different family-group taxa.

7. To illustrate the prevalence of the name PSEUDOCOCCIDAE, the following references are provided: Ben-Dov, 1994; Miller et al., 2002; Bonnett & Hewitt, 2005; Gonzalez-Hernandez et al., 2005; Kondo et al., 2005; Wakgari & Giliomee, 2005; Hogendorp et al., 2006; Kaydan et al., 2006; Martinez-Ferrer et al., 2006; Williams & Moghaddam, 2007. A list of 499 additional references demonstrating the universal usage of the name PSEUDOCOCCIDAE is held by the Commission Secretariat and is available upon request.

8. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary power to rule that the family-group name PSEUDOCOCCINI Cockerell, 1905 and other family-group names based on *Pseudococcus* Westwood, 1840 are to be given precedence over SPHAEROCOCCINI Cockerell, 1899 and other family-group names based on *Sphaerococcus* Maskell, 1892 whenever their type genera are placed in the same family-group taxon;
- (2) to place on the Official List of Generic Names in Zoology the name *Sphaerococcus* Maskell, 1892 (gender: masculine), type species by monotypy *S. casuarinae* Maskell, 1892;
- (3) to place on the Official List of Family-Group Names in Zoology the names:
  - (a) PSEUDOCOCCINI Cockerell, 1905 (type genus *Pseudococcus* Westwood, 1840), with the endorsement that it and other family-group names based on *Pseudococcus* are to be given precedence over SPHAEROCOCCINI Cockerell, 1899 and other family-group names based on *Sphaerococcus* Maskell, 1892 whenever their type genera are placed in the same family-group taxon;
  - (b) SPHAEROCOCCINI Cockerell, 1899 (type genus *Sphaerococcus* Maskell, 1892), with the endorsement that it and other family-group names based on *Sphaerococcus* are not to be given priority over PSEUDOCOCCINI Cockerell, 1905 and other family-group names based on *Pseudococcus* Westwood, 1840 whenever their type genera are placed in the same family-group taxon.

## References

- Ben-Dov, Y. 1994. *A Systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data and geographic distribution, host plants, biology and economic importance*. 686 pp. Intercept Ltd, Andover.
- Bonnett, G.D. & Hewitt, M.L. 2005. Numbers of pink sugarcane mealy bug, *Saccharicoccus sacchari* (Cockerell) (Hemiptera: Pseudococcidae) differ within seasons and among regions and stages of the sugarcane crop cycle. *Australian Journal of Entomology*, **44**: 304–309.
- Cockerell, T.D.A. 1899. Article VII. – First supplement to the check-list of the Coccidae. *Bulletin of the Illinois State Laboratory of Natural History*, **5**: 389–398.
- Cockerell, T.D.A. 1905. Tables for the identification of Rocky Mountain Coccidae (Scale insects and mealybugs). *Colorado University Studies*, **2**: 189–203.
- Gonzalez-Hernandez, H., Pandey, R.R. & Johnson, M.W. 2005. Biological characteristics of adult *Anagyrus ananatis* Gahan (Hymenoptera: Encyrtidae), a parasitoid of *Dysmicoccus brevipes* (Cockerell) (Hemiptera: Pseudococcidae). *Biological Control*, **35**: 93–103.

- Hogendorp, B.K., Cloy, R.A. & Swiader, J.M.** 2006. Effect of nitrogen fertility on reproduction and development of citrus mealybug, *Planococcus citri* Risso (Homoptera: Pseudococcidae), feeding on two colors of coleus, *Solenostemon scutellarioides* L. Codd. *Environmental Entomology*, **35**: 201–211.
- Kaydan, M.B., Kilincer, N., Uygun, N., Japoshvilli, G. and Gaimari, S.** 2006. Parasitoids and predators of Pseudococcidae (Hemiptera: Coccoidea) in Ankara, Turkey. *Phytoparasitica*, **34**: 331–337.
- Kondo, T., Gullan, P.J., Ventura, J.A. & Culik, M.P.** 2005. Taxonomy and biology of the mealybug genus *Plotococcus* Miller & Deno (Hemiptera: Pseudococcidae) in Brazil, with descriptions of two new species. *Studies on Neotropical Fauna and Environment*, **40**: 213–227.
- Lobdell, G. H.** 1930. Twelve new mealybugs from Mississippi (Homoptera: Coccoidea). *Annals of the Entomological Society of America*, **23**: 209–236.
- Martinez-Ferrer, M.T., Ripolles, J.L. & Garcia-Mari, F.** 2006. Enumerative and binomial sampling plans for citrus mealybug (Homoptera: Pseudococcidae) in citrus groves. *Journal of Economic Entomology*, **99**: 993–1001.
- Maskell, W.M.** 1892 (1891). Further coccid notes: with descriptions of new species, and remarks on coccids from New Zealand, Australia and elsewhere. *Transactions and Proceedings of the New Zealand Institute*, **24**: 1–64.
- Miller, D.R.** 1974. *Dactylopius* Costa, 1835 and *Pseudococcus* Westwood, 1840 (Insecta Homoptera): proposed designation of type-species under plenary powers with proposed suppression of *Diaprosteci* Costa, 1828. *Bulletin of Zoological Nomenclature*, **31**: 146–153.
- Miller, D.R., Miller, G.L. & Watson, G.W.**, 2002. Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to U.S. agriculture. *Proceedings of the Entomological Society of Washington*, **104**: 825–836.
- Targioni-Tozzetti, M.H.** 1867. Sur la cire qu'on peut obtenir de la cochenille du figuier (*Coccus caricae*) auct. *Compte Rendu de l'Académie des Sciences. Paris*, **45**: 246–247.
- Wakgari, W.M. & Giliomee, J.H.** 2005. Description of adult and immature females of six mealybug species (Hemiptera: Pseudococcidae) found on citrus in South Africa. *African Entomology*, **13**: 281–332.
- Westwood, J.O.** 1840. *An introduction to the modern classification of insects; founded on the natural habits and corresponding organization of different families*, vol. 2. 587 pp. Longman, Orme, Brown, Green and Longmans, London.
- Williams, D.J. & Moghaddam, M.** 2007. A new species of *Rhodania* Goux (Hemiptera: Coccoidea, Pseudococcidae) and new records of two other mealybug species from Iran. *Zootaxa*, **1398**: 37–44.

Acknowledgement of receipt of this application was published in BZN **64**: 138.

---

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).